

The CRAFFT Screening Questions

Please answer all questions honestly; your answers will be kept confidential.

Part A

During the PAST 12 MONTHS, did you:

- | | | | |
|---|--------------------------|--|--------------------------|
| | No | | Yes |
| 1. Drink any <u>alcohol</u> (more than a few sips)? | <input type="checkbox"/> | | <input type="checkbox"/> |
| 2. Smoke any <u>marijuana or hashish</u> ? | <input type="checkbox"/> | | <input type="checkbox"/> |
| 3. Use <u>anything else</u> to <u>get high</u> ? | <input type="checkbox"/> | | <input type="checkbox"/> |

“anything else” includes illegal drugs, over the counter and prescription drugs, and things that you sniff or “huff”

If you answered NO to ALL (A1, A2, A3) answer **only B1** below, then STOP.

If you answered YES to ANY (A1 to A3), answer **B1 to B6** below.

Part B

- | | | |
|--|--------------------------|--------------------------|
| | No | Yes |
| 1. Have you ever ridden in a CAR driven by someone (including yourself) who was “high” or had been using alcohol or drugs? | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. Do you ever use alcohol or drugs to RELAX, feel better about yourself, or fit in? | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. Do you ever use alcohol or drugs while you are by yourself, or ALONE? | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. Do you ever FORGET things you did while using alcohol or drugs? | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. Do your FAMILY or FRIENDS ever tell you that you should cut down on your drinking or drug use? | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. Have you ever gotten into TROUBLE while you were using alcohol or drugs? | <input type="checkbox"/> | <input type="checkbox"/> |

CONFIDENTIALITY NOTICE:

The information on this page may be protected by special federal confidentiality rules (42 CFR Part 2), which prohibit disclosure of this information unless authorized by specific written consent. A general authorization for release of medical information is NOT sufficient.


© Children’s Hospital Boston, 2009.

Reproduced with permission from the Center for Adolescent Substance Abuse Research, CeASAR, Children’s Hospital Boston. CRAFFT Reproduction produced with support from the Massachusetts Behavioral Health Partnership.

SCORING INSTRUCTIONS: FOR CLINIC STAFF USE ONLY

CRAFFT Scoring: Each “yes” response in **Part B** scores 1 point.
A total score of 2 or higher is a positive screen, indicating a need for additional assessment.

Probability of Substance Abuse/Dependence Diagnosis Based on CRAFFT Score^{1,2}



DSM-IV Diagnostic Criteria³ (Abbreviated)

Substance Abuse (1 or more of the following):

- Use causes failure to fulfill obligations at work, school, or home
- Recurrent use in hazardous situations (e.g. driving)
- Recurrent legal problems
- Continued use despite recurrent problems

Substance Dependence (3 or more of the following):

- Tolerance
- Withdrawal
- Substance taken in larger amount or over longer period of time than planned
- Unsuccessful efforts to cut down or quit
- Great deal of time spent to obtain substance or recover from effect
- Important activities given up because of substance
- Continued use despite harmful consequences

© Children’s Hospital Boston, 2009. This form may be reproduced in its exact form for use in clinical settings, courtesy of the Center for Adolescent Substance Abuse Research, Children’s Hospital Boston, 300 Longwood Ave, Boston, MA 02115, U.S.A., (617) 355-5433, www.ceasar.org.

References:

1. Knight JR, Shrier LA, Bravender TD, Farrell M, Vander Bilt J, Shaffer HJ. A new brief screen for adolescent substance abuse. Arch Pediatr Adolesc Med 1999;153(6):591-6.
2. Knight JR, Sherritt L, Shrier LA, Harris SK, Chang G. Validity of the CRAFFT substance abuse screening test among adolescent clinic patients. Arch Pediatr Adolesc Med 2002;156(6):607-14.
3. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, fourth edition, text revision. Washington DC, American Psychiatric Association, 2000.